


Pneumococcal Vaccine Timing For Adults

DO NOT administer PCV13 and PPSV23 at the same visit.

Age 65 Years or Older

- If PCV13 was given before age 65 years, no additional PCV13 is needed.



Age 19-64 Years With Underlying Condition(s)

- Prior doses count toward doses recommended below and do not need to be repeated.
- If PPSV23 given previously – wait one year before giving PCV13
 - for group B, wait at least five years before giving a second dose of PPSV23
- No more than two doses of PPSV23 recommended before 65th birthday and one dose thereafter.

A. Smoker, Long-term facility resident, or Chronic conditions:

- heart disease (excluding hypertension)
- lung disease (including asthma)
- liver disease (including cirrhosis)
- diabetes
- alcoholism

PPSV 23

B. Immunocompromised (including HIV infection), Chronic renal failure, Nephrotic syndrome, or Asplenia

PCV 13

8 weeks

PPSV 23

5 years

PPSV 23

C. CSF leaks or Cochlear implants

PCV 13

8 weeks

PPSV 23



For further details, see: www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/pneumo.html

Reproduced with permission from the California Department of Public Health, Immunization Branch. Development was supported by Grant H23/CCH922507 from the Centers for Disease Control and Prevention.

This material was adapted by HealthInsight, the Medicare Quality Innovation Network (QIN) Quality Improvement Organization for Nevada, New Mexico, Oregon and Utah, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. The contents presented do not necessarily reflect CMS policy. 11SOW-F1-16-02-NM